ZWANGER-PESIRI RADIOLOGY 631-444-5544

MEDICAL ASSOCIATES

						/ /
PATIENT LAST NAME		PATIENT FULL FIRST NAM	IE	T	ODAY'S DATE	DATE OF BIRTH
CLINICAL INDICATION	IS/SIGNS/SYMPTOMS	(NOT RULE/OUT): ——			. [
ICD-10:						
PHYSICIAN SIGNA	ATURE (REQUIRED)	775 Park Avenue, Suite 145, Huntington, NY 11743 T: (631) 261-4445 F: (631) 944-3018				
PATIENTS: Call to make an appointment of take a cell phone photo of this form and text or email it to rx@zprad.com						
= MOI		СТ				
MRI		□ With Contrast □ Without Contrast □ With & Without Contrast		Contrast	■ Mammography □ Please schedule breast sonogram appointment if	
☐ 3T Wide-Bore ☐ 1.5T Wide-Bore ☐ 1.2 Open-Sided ☐ Either 3T or 1.5T Wide-Bore		☐ Oral Contrast Only ☐ IV Contrast Only ☐ Oral & IV Contrast		ntrast	needed based on the mammogram. Screening With 3D Tomosynthesis	
☐ With & without contrast ☐ No contrast		CT Angiography □ Coronary artery CTA	Spine □ Cervical □ Thoracic Specify levels		(no palpable finding or symptoms) □ Bilateral □ Right □ Left □ Screening (no palpable finding or symptoms) □ Bilateral □ Right □ Left	
☐ With I.V. sedation		with calcium scoring (needs contrast)				
Neuro/ENT/Spine	MRA	☐ Chest CTA/PE ☐ Calcium scoring only	Lumbar			osynthesis-Must select reason(s)
☐ Brain☐ Orbits	☐ Carotid MRA☐ Intracran/circle of Willis	☐ CT angiogram (needs contrast) ☐ Intracranial	☐ Sacrum/coccyx		☐ Bilateral ☐ Right ☐ Diagnostic - Must select	t □Left
☐ Pituitary ☐ IAC	☐ Intracran/MR venogram ☐ MR venogram	☐ Carotid ☐ Aortic arch/thoracic	Body ☐ Stone hunt		☐ Bilateral ☐ Right	
☐ Cervical spine ☐ Thoracic spine	Specify	aorta	☐ Hematuria ☐ Chest only		Reasons:	views
☐ Lumbar spine ☐ Sacrum/coccyx	□Carotid	☐ Lower extremity run off	☐ Soft tissues neck/c	hest/	☐ Short term follow up ☐ New lump, mass or th	
☐ CSF Flow	☐ Aortic arch ☐ Abdominal aorta only	Neuro/ENT □ Brain	abdomen/pelvis ☐ Soft tissues neck or	only	☐ Old lump or mass incl	reased in size
☐ DTI ☐ Perfusion	☐ Renal arteries ☐ Mesenteric arteries	☐ Orbits ☐ Temporal bones	☐ Chest/abdomen/pel ☐ Abdomen/pelvis	pelvis	☐ New inverted nipple	
☐ MR spectroscopy ☐ TMJ	☐ Aorta/lower extremities	☐ Paranasal sinuses ☐ Soft tissues neck	☐ Enterography ☐ Abdomen only		☐ Skin changes (dimplir increase in breast size)	ng, redness or abnormal
☐ Soft tissue neck/parotid	Chest & Body □Chest	Musculoskeletal □ Joint	☐ Pelvis only		☐ Lymphadenopathy ☐ Current use of Tamoxi	ifen, Femara or Arimidex
Orthopedic ☐ Shoulder ☐ R ☐ L	☐ Breast MRI	Specify	☐ Triple phase liver		■Ultrasound	
□Upper arm □R □L	☐ Cardiac MRI☐ Function☐ Viability☐	Specify	☐ Other		□ Breast	Vascular
☐ Elbow ☐ R ☐ L ☐ Forearm ☐ R ☐ L	☐ Mediastinum ☐ Brachial plexus	■ Nuclear Madia	••••		□ Bilateral □ R □ L □ Thyroid	☐ Carotid doppler☐ Venous doppler
□Wrist □R□L □Hand □R□L	☐ Clavicle/sc joint☐ Scapula	■ Nuclear Medic	Thyroid		☐ Scrotal/testicular ☐ Transrectal prostate	☐ Lower extremity
☐ Finger ☐ R ☐ L Specify	☐ Sternum ☐ Thoracic outlet	☐ Add SPECT if needed	□ Uptake & scan □ I-131 treatment	,	☐ Pelvis (GYN)	☐ R ☐ L ☐ Bilateral ☐ Upper extremity
☐ Pelvis ☐ R ☐ L	□Abdomen	☐ Whole body	Dose		☐ Transabdominal ☐ Transvaginal	☐ R ☐ L ☐ Bilateral ☐ Arterial doppler
☐ Hip ☐ R ☐ L ☐ L ☐ L ☐ L	Specify	☐ 3 phase Region	□With cholecysto □Renal	okinin	☐ Transabdominal / Transvaginal	☐ Lower extremity
☐ Knee ☐ R ☐ L ☐ Lower leg ☐ R ☐ L	☐ Dynamic pelvis/ MR defogram	☐ Cardiac	☐ With lasix washo ☐ DTPA	out	☐ Hysterosonogram ☐ Obstetrical	☐ R ☐ L ☐ Bilateral ☐ Upper extremity
☐ Ankle ☐ R ☐ L ☐ Foot ☐ R ☐ L	☐ Prostate ☐ Enterography	☐ Myocardial perfusion stress	☐ Parathyroid ☐ Gastric emptying		□ Abdomen	☐ R ☐ L ☐ Bilateral☐ Renal arterial doppler
☐Toe ☐R ☐L ☐ Cartilage mapping	□MRCP	study □With treadmill/	☐ Other		☐ Aorta only ☐ Retroperitoneum	Tenar arteriar doppier
☐ MR arthrogram	☐ Rectal MRI	exercise □With pharm. agent			(Renal/Bladder)	
Specify Other		☐ MUGA (gated blood pool)			Other	
					■ Echocardiogram	
■ DXA Bone Density		■ PET/CT ■ Add CT intravenous contrast if needed			■ Interventional Biopsy □Thyroid □Lung □Liver	
		PET/CT Auth#:			☐ US Breast FNA Specify Region	
■ Fluoroscopy		☐ 78608 Brain PET ☐ 78815 Base of skull to mid thigh ☐ 78816 Top of head to toes (melanoma protocol)			Specify Region Stereotactic Biopsy (includes post procedure mammo)	
☐ Lap band ☐ Hysterosalpingogram				ol)	Specify Region □ Perform targeted US first, if lesion identified, biopsy under US	
□ Other:		□ Other:			☐ MRI Breast Biopsy 1 Specify Region ☐ Perform targeted US first, if lesion identified, biopsy under US	
■ Digital X-RAY Patients can print registration forms online □ Skull □ C spine □ C sp						
□ Orbits □ T spine □ F/U abdomen □ Ribs □ Humerus □ R □ L □ Hand □ R □ L □ Knee □ R □ L □ Toes □ R □ L						□L □Toes □R □L
□ Facial bones □ □ L spine □ □ KUB abdomen □ □ Elbow □ R □ L □ Fingers □ R □ L □ □ Tibia/fibula □ R □ L □ Other: □ Nasal bones □ Sacrum □ Pelvis □ Forearm □ R □ L □ Hips □ R □ L □ Ankle □ R □ L □ Other:						

EXAM PREPARATIONS



- · Continue taking any prescription medications, which may be taken with a few sips of water prior to exam.
- Wear comfortable, loose clothing. Do not wear jewelry.
 Please be sure you have your prescription from the doctor, as well as your insurance card when you arrive for your appointment.

MRI/MRA

★ If you are receiving IV CONTRAST for your exam, have nothing to eat 1 hour prior to your exam time. You may drink clear liquids (example: water, ginger ale, apple juice).

This exam may not be performed if you have a cardiac pacemaker, cerebral aneurysm clips or a cochlear ear implant. If you are a sheet metal worker or have ever had **metal fragments in your eye(s)**, an orbit x-ray may be taken prior to your MRI exam. Wear comfortable loose fitting clothes, such as a sweatsuit. Be sure there are no metal zippers, snaps or buckles. Do not wear earrings, hairpins or jewelry. Do not apply eye shadow or mascara.

CT SCAN

- ★ If you have a history of asthma, an allergy to iodine, or are currently taking medication for diabetes, please notify our staff.
- ★ If you are receiving IV CONTRAST or ORAL CONTRAST for your exam, have nothing to eat 1 hour prior to your exam time. You may drink clear liquids (example: water, ginger ale, apple juice). If you are receiving OMNIPAQUE ORAL CONTRAST, refer to the OMNIPAQUE ORAL CONTRAST PREP section below. If you are receiving **REDI-CAT ORAL** CONTRAST, please ask your Zwanger-Pesiri representative.

Abdomen and Pelvis with or without contrast - Nothing to eat or drink 1 hour prior to your exam time.

All other exams with no contrast - No preparation necessary.

CT Angiography - Follow instructions given at the time of scheduling.

OMNIPAQUE ORAL CONTRAST PREP FOR CT SCAN OF ABDOMEN AND PELVIS

★ DO NOT INGEST IF YOU HAVE A IODINE ALLERGY

- · Have nothing to eat 1 hour prior to your exam time.
- You may drink clear liquids (example: water, ginger ale, apple juice).
- Begin drinking the OMNIPAQUE prep 1 hour & 45 minutes before your exam. Finish in 15 minutes.
- Do not empty your bladder until after your exam.
- Pour HALF the contents of the OMNIPAQUE bottle into the 32 oz. cup given at the office.
- 2 Fill the cup with water to approximately 1/2 inch from the top of the cup (approximately 30 oz).
- Stir well and drink.
- ② Discard the remaining contrast, cup, contrast bottle and straw after use.

PET/CT SCAN

Call your local Zwanger-Pesiri office to schedule an appointment and/or for exam preparations. MRI/PET: In addition, follow second paragraph of MRI preparations.

3D & 2D DIGITAL MAMMOGRAPHY

Do not apply lotions, deodorant, perfume or powder on the day of the exam. Wear comfortable two piece clothing. Bring previous mammography studies for comparison.

DXA BONE DENSITOMETRY

No calcium supplements or multi-vitamins 24 hours prior to exam.

SPECT NUCLEAR MEDICINE

Call your local Zwanger-Pesiri office to schedule an appointment.

Thyroid Scan - Discontinue all thyroid medications and vitamins with iodine for at least 10 days prior to the exam. Continue any beta blockers as prescribed. Please advise us at the time of scheduling if you have received intravenous CT contrast within the last 6 weeks.

Bone Scan - No preparation required.

Liver Scan - No preparation required.

ULTRASOUND

Aortic/Abdominal - Nothing to eat drink, chew or smoke for six hours prior to your exam.

Pelvic/Obstetrical - A full bladder is necessary for the exam. Have breakfast and/or lunch. Women: drink at least 32 oz. of water/Men: at least 16 oz. of water, finishing 1 hour prior to exam. Do not empty your bladder.

Prostate - Take a fleet enema at least one hour prior to the exam. Nothing to eat or drink after fleet enema.

Breast/Scrotal/Thyroid - No preparation required.

Extremity Doppler - No preparation required.

Renal Arterial Study - Nothing to eat, drink, chew or smoke for six hours prior to your exam. In addition, consult your physician before taking gas-X one hour before the exam.

Renal - 16 oz. glass of water one hour prior to study. Do not void.

